

Distinguished bases of exceptional modules.

Claus Michael Ringel

An indecomposable representation M of a quiver $Q = (Q_0, Q_1)$ is said to be exceptional provided $\text{Ext}^1(M, M) = 0$. And it is called a tree module provided one can choose a set $\mathcal{B}$ of bases of the vector spaces M_x ($x \in Q_0$) such that the coefficient quiver $\Gamma(M, \mathcal{B})$ is a tree quiver; we call $\mathcal{B}$ a tree basis of M . It is known that exceptional modules are tree modules. Now a tree module usually has many tree bases and the corresponding coefficient quivers may look quite differently. The aim of this note is to introduce a class of exceptional modules which have a distinguished tree basis, the “radiation modules”. For a Dynkin quiver Q , nearly all indecomposable representations turn out to be radiation modules, the only exception is the maximal indecomposable module in case $\mathbb{E}_8$. Also, the exceptional representation of the generalized Kronecker quivers are given by radiation modules. Consequently, with the help of Schofield induction one can display all the exceptional modules of an arbitrary quiver in a nice way.

Let $Q = (Q_0, Q_1)$ be a locally finite quiver. We will consider finite-dimensional representations of Q (thus kQ -modules, where kQ is the path algebra of Q). The quivers we are interested will be directed without infinite paths. An indecomposable representation M of Q is said to be *exceptional* provided $\text{Ext}^1(M, M) = 0$. And it is called a *tree module* provided one can choose a set $\mathcal{B}$ of bases of the vector spaces M_x ($x \in Q_0$) such that the coefficient quiver $\Gamma(M, \mathcal{B})$ (see [R2]) is a tree quiver; we call $\mathcal{B}$ a *tree basis* of M . It is known that exceptional modules are tree modules. But even if we know that a module M is a tree module, it often seems to be difficult to find directly a tree basis. The aim of this note is to provide for the exceptional modules an algorithm for obtaining a tree basis. The first modules which we will consider are what we call the radiation modules, see section 2. They are inductively defined, in any step one constructs an exceptional module with a thin vertex. Note that we say that a vertex x is thin for the module M provided the vector space M_x is one-dimensional. In section 1 we are analyzing the exceptional modules with a thin vertex.

As we will see in section 3, nearly all indecomposable representation of a Dynkin quiver are radiation modules, the only exception is the maximal indecomposable module in case $\mathbb{E}_8$. The radiation modules which are our main concern are the preprojective and preinjective representation of a tree without leafs with bipartite orientation (a leaf of a tree is a vertex with a single neighbor). These modules are studied in sections 4. Typical examples of trees without leafs are the n -regular trees with $n \geq 2$, note that they occur when we consider the universal covering of a generalized Kronecker quiver. Thus, using covering theory, one obtains a distinguished tree basis for any exceptional representation of a generalized Kronecker module. And with the help of Schofield induction we then get a nice tree basis for any exceptional module, see the outline in the last section 5.

1. Exceptional representations M with a thin vertex.

If M is a representation of Q and $\dim M_x = 1$ for some vertex x , we say that x is a *thin* vertex for M .

Let Q be a quiver with underlying graph $\overline{Q}$ being a tree. Let x be a vertex of Q and let Q^x be obtained from Q by deleting the vertex x and all the arrows involving x .

We say that a sequence of modules $N(1), \dots, N(t)$ is an *orthogonal exceptional family*, provided all the modules $N(i)$ are exceptional, and $\text{Hom}(N(i), N(j)) = 0 = \text{Ext}^1(N(i), N(j))$ for all $i \neq j$.


Proposition 1. *Let M be an exceptional representation of Q with thin vertex x . Then the restriction M' of M to Q^x is of the form $M' = \bigoplus_{i=1}^t N(i)^{n(i)}$, where $N(1), \dots, N(t)$ is an orthogonal exceptional family, and all $n(i) > 0$, such that for any index i , there is a neighbor $y(i)$ of x with $\dim N(i)_{y(i)} = n(i)$ and this is the only neighbor y of x with $\dim N(i)_y \neq 0$.*

Conversely, let $N(1), \dots, N(t)$ be an orthogonal exceptional family with $N(i)_x = 0$ and assume that for any index i , there is a neighbor $y(i)$ of x with $\dim N(i)_{y(i)} = n(i) > 0$. Then there is an exceptional representation M of Q , unique up to isomorphism, with thin vertex x such that the restriction of M to Q^x is $M' = \bigoplus_{i=1}^t N(i)^{n(i)}$.

Proof: First, let us assume that modules $N(1), \dots, N(t)$ are given which are pairwise orthogonal and exceptional, with $N(i)_x = 0$, and that for any index i , there is a neighbor $y(i)$ of x with $\dim N(i)_{y(i)} = n(i) > 0$. We can assume that there are arrows $x(i) \rightarrow x$ for $1 \leq i \leq s$ and arrows $x \rightarrow y(i)$ for $s+1 \leq i \leq t$. Since $N(i)_x = 0$, we see that the modules $S(x), N(1), \dots, N(t)$ are orthogonal and that the sequence

$$N(1), \dots, N(s), S(x), N(s+1), \dots, N(t)$$

is exceptional. Note that the Ext-quiver of this set of modules is as follows:



with $n(i)$ arrows between $N(i)$ and $S(x)$, for $1 \leq i \leq t$.

Now such a quiver has the following real root:

$$\begin{array}{ccccccc} n(1) & n(2) & \cdots & n(s) & & & \\ & & & & 1 & & \\ n(s+1) & n(s+2) & \cdots & n(t) & & & \end{array}$$

and the process of simplification shows that the category of representations of Q which have a filtration with factors in $\{S(x), N(1), \dots, N(t)\}$ has precisely one indecomposable

object with one factor $S(x)$ and $n(i)$ factors $N(i)$, for $1 \leq i \leq t$, and that this is an exceptional module.

For the converse, we use the following lemma.

Lemma 1. *Let X' be a submodule of X and $X'' = X/X'$. We assume that $\text{Ext}^1(X, X) = 0$ and $\text{Hom}(X', X'') = 0$. Then $\text{Ext}^1(X', X') = 0$ and $\text{Ext}^1(X'', X'') = 0$.*

Proof: The inclusion map $X' \rightarrow X$ yields an epimorphism $\text{Ext}^1(X, X) \rightarrow \text{Ext}^1(X', X)$, thus $\text{Ext}^1(X', X) = 0$. Applying $\text{Hom}(X', -)$ to the exact sequence $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$ yields an exact sequence

$$\text{Hom}(X', X'') \rightarrow \text{Ext}^1(X', X') \rightarrow \text{Ext}^1(X', X).$$

Since the end terms are zero, also the middle term is zero. This shows that $\text{Ext}^1(X', X') = 0$. By duality, also $\text{Ext}^1(X'', X'') = 0$.

Assume now that M is an exceptional representation of Q such that $\dim M_x = 1$ and denote by M' the restriction of M to Q^x . Let r be the number of neighbors y of x such that $M_y \neq 0$.

We first consider the case $r = 1$, thus there is a unique neighbor y of x such that $M_y \neq 0$. We may assume that there is an arrow $x \rightarrow y$ (the dual case of an arrow $y \rightarrow x$ follows by duality). Let M' be the restriction of M to Q^x . Then M' is a submodule of M with factor module $M/M' = S(x)$. Since we have $\text{Hom}(M', M/M') = 0$, lemma 1 asserts that $\text{Ext}^1(M', M') = 0$. Write $M' = \bigoplus N(i)^{n(i)}$ with pairwise non-isomorphic indecomposable modules $N(i)$, then $\text{Ext}^1(N(i), N(j)) = 0$ for all i, j ; in particular, the modules $N(i)$ are exceptional. Also, $\dim \text{Ext}^1(S(x), N(i)) = \dim N(i)_y = n(i)$ for all i , thus $\dim \text{Ext}^1(S(x), M') = \sum_i n(i) \dim \text{Ext}^1(S(x), N(i)) = \sum_i n(i)^2$. If we apply $\text{Hom}(S(x), -)$ to the exact sequence

$$(*) \quad 0 \rightarrow M' \rightarrow M \rightarrow S(x) \rightarrow 0,$$

we obtain an exact sequence

$$\text{Hom}(S(x), M) \rightarrow \text{Hom}(S(x), S(x)) \rightarrow \text{Ext}^1(S(x), M') \rightarrow \text{Ext}^1(S(x), M) \rightarrow \text{Ext}^1(S(x), S(x)).$$

Since there are no loops in Q , the last term is zero. Also, the first term is zero, since otherwise $M = S(x)$ and $r = 0$. It follows that

$$\dim \text{Ext}^1(S(x), M') = -1 + \sum n(i)^2.$$

Now we apply $\text{Hom}(-, M)$ to $(*)$ and obtain an exact sequence

$$\text{Hom}(S(x), M) \rightarrow \text{Hom}(M, M) \rightarrow \text{Hom}(M', M) \rightarrow \text{Ext}^1(S(x), M) \rightarrow \text{Ext}^1(M, M).$$

As we have seen, the first term is zero, and by assumption also the last term is zero. Since $\text{Hom}(M, M) = k$, we see that

$$\dim \text{Hom}(M', M) = 1 + \dim \text{Ext}^1(S(x), M') = \sum n(i)^2.$$

But any homomorphism $M' \rightarrow M$ maps into M' , thus $\dim \operatorname{Hom}(M', M) = \dim \operatorname{Hom}(M', M')$. But $\operatorname{Hom}(M', M') = D \oplus E$, with subspaces

$$D = \bigoplus_i \operatorname{Hom}(N(i)^{n(i)}, N(i)^{n(i)}), \quad E = \bigoplus_{i \neq j} \operatorname{Hom}(N(i)^{n(i)}, N(j)^{n(j)}).$$


Since $\dim D = \sum_i n(i)^z$, it follows that $E = 0$, thus $\operatorname{Hom}(N(i), N(j)) = 0$ for all $i \neq j$. This completes the proof in case $r = 1$.

Now assume that there are $r > 1$ neighbors y of x with $M_y \neq 0$, say $y = y_1, \dots, y_r$. Let $Q^x(i)$ be the connected component of Q^x which contains the vertex y_i . Decompose $M' = \bigoplus_{i=1}^r M'(i)$ such that $M'(i)$ has support in $Q^x(i)$. Of course, this implies that $\operatorname{Hom}(M'(i), M'(j)) = 0$ for $i \neq j$. We also have $\operatorname{Ext}^1(M'(i), M'(j)) = 0$ for $i \neq j$, since there are no arrows from vertices of $Q^x(i)$ to vertices of $Q^x(j)$.

Let $Q(i)$ be obtained from $Q^x(i)$ by adding the vertex x as well as the arrow which connects x and y , and let $M(i)$ be the restriction of M to $Q(i)$. We claim that $M(i)$ is an exceptional module. Clearly, $M(i)$ is indecomposable: any non-trivial direct decomposition of $M(i)$ yields a corresponding direct decomposition of M , since $\dim M_x = 1$. Also, using lemma 1 and induction, one sees easily that $\operatorname{Ext}^1(M(i), M(i)) = 0$.


Now $\dim M(i)_x = 1$ and there is a unique neighbor y of x with $M(i)_y \neq 0$. Thus looking at the module $M(i)$ and the vertex x , we are in the case $r = 1$ of the proposition, thus we know already that $M'(i)$ decomposes as required. But the decompositions of the modules $M'(i)$ combine to the required decomposition of M' , since $\operatorname{Hom}(M'(i), M'(j)) = 0 = \operatorname{Ext}^1(M'(i), M'(j))$ for all $i \neq j$.

Example 1. Consider the following exceptional representation M , encircled is a vertex x with $\dim M_x = 1$, note that x has a unique neighbor y . The lower line exhibits the dimension vectors of the modules $N(i)$, here the vertex y is encircled by a dotted line.



We mainly will be interested in the case that $n(i) = 1$ for all i , thus in the case that M' decomposes as the direct sum of pairwise non-isomorphic indecomposable modules $N(i)$. Here is an example with $n(i) > 1$ for some index i .

Example 2. An exceptional module M with a thin vertex x such that the restriction M' of M to Q^x is of the form $N(1) \oplus N(2)^2$ for an exceptional orthogonal family $N(1), N(2)$.



2. Radiation modules.

Let us assume that we deal with a quiver Q which is a tree. We consider pairs (M, x) where M is an indecomposable module and x is a vertex with $\dim M_x = 1$ (called the *origin* of M or better of the pair (M, x)). We define the class of *radiation modules* as well as the corresponding *radiation quivers* inductively as follows. First of all, for any vertex x , the pair $(S(x), x)$ is a radiation module, its radiation quiver $R(S(x), x)$ is the quiver with a single vertex (and no arrow), the arrow being labeled x .

Second, if M is an indecomposable module of length at least 2 and x is vertex with $\dim M_x = 1$, then (M, x) is a radiation module proved the restriction of M to the quiver obtained from Q by deleting the vertex x is the direct sum $\bigoplus N(i)$ of indecomposable modules $N(i)$ such that for any index i there is a neighbor $y(i)$ of x with $(N(i), y(i))$ being a radiation module. The radiation quiver $R(M, x)$ of (M, x) is obtained from the disjoint union of the radiation quivers $R(N(i), y(i))$ by adding a vertex with label x and for every index i we connect x and $y(i) \in R(N(i), y(i))$ by an arrow which points in the same direction as the arrow in Q between x and $y(i)$, and we use this arrow of Q also as the label of the arrow in $R(M, x)$. Note that the labels of the vertices and the arrows of $R(M, x)$ provide a quiver homomorphism $R(M, x) \rightarrow Q$.


Proposition 2. *If (M, x) is a radiation module, then M is exceptional and $R(M, x)$ is a coefficient quiver of M . Radiation quivers are always trees.*

Proof. The last assertion is straight-forward: Namely, assume that there is given a finite set of trees $T(i)$ and for every i a vertex $y(i)$ in $T(i)$. If we take the disjoint union of the graphs $T(i)$, an additional vertex x and connect x with the vertices $y(i)$ by a single edge, then we obtain again a tree.

The first assertion follows from the proposition 1.

It remains to show that $R(M, x)$ is a coefficient quiver of M . Let b be a non-zero element of M_x (recall that M_x is one-dimensional). We define a basis $\mathcal{B}(M, b)$ of M as follows: Assume that the restriction of M to the quiver obtained by deleting the vertex x is $\bigoplus N(i)$ with indecomposable representations $N(i)$. For every index i , there is a vertex $y(i)$ which is a neighbor of x such that $(N(i), y(i))$ is a radiation module. Since $x, y(i)$ are neighbors, there is an arrow say α which connects them. If $\alpha: x \rightarrow y(i)$, then let $\mathcal{B}(i) = \mathcal{B}(N(i), \alpha(b))$. If $\alpha: y(i) \rightarrow x$, then let $\mathcal{B}(i) = \mathcal{B}(N(i), \alpha^{-1}(b))$. Finally, let $\mathcal{B}(M, B)$ be the set containing the element b as well as all the elements in the (disjoint!) sets $\mathcal{B}(i)$. By induction we see that $\mathcal{B}(M, B)$ is a basis of M and the corresponding coefficient quiver is just $R(M, x)$.

Example of a tree basis which is not a radiation basis. We deal with the quiver Q of type $\mathbb{D}_5$ with subspace orientation. We denote by Γ a coefficient quiver of the maximal indecomposable module M :



There are three vertices x with $\dim M_x = 1$, namely the leaves of the quiver Q . Removing any such vertex x , the module $M|(Q \setminus \{x\})$ is the direct sum of two indecomposables, thus in $\mathcal{B}(M, x)$, there are two arrows starting in x . But note that dealing with Γ , only one arrow starts at any leaf.

Properties of a radiation quiver. Let us start with a general (and well-known) property of exceptional modules:

Lemma 2. *Let M be a representation of the quiver Q with $\text{Ext}^1(M, M) = 0$. Then for every arrow α in Q , the linear map M_α is injective or surjective.*

Proof: Assume that M_α is not bijective for $\alpha: x \rightarrow y$. Write $M_x = M'_x \oplus M''_x$ with M'_x the kernel of M_α and $M_y = M'_y \oplus M''_y$ with M'_y the cokernel of M_α . By assumption M'_x and M''_y are non-zero, thus there is a non-zero linear map $N: M_x \rightarrow M_y$ such that $N(M'_x) \subseteq M''_y$ and $N(M''_x) = 0$.

In order to construct a non-trivial self-extension of M , start with the direct sum $M \oplus M$ and replace the map $(M \oplus M)_\alpha = \begin{bmatrix} M_\alpha & 0 \\ 0 & M_\alpha \end{bmatrix}$ by the map $\begin{bmatrix} M_\alpha & N \\ 0 & M_\alpha \end{bmatrix}$. The properties of N show that the rank of this map is larger than the rank of $(M \oplus M)_\alpha$, thus the new representation cannot be isomorphic to $M \oplus M$, and obviously, it is an extension of M by itself.

Now let (M, x) be a radiation module. We consider a path

$$x \text{ --- } \cdots \text{ --- } y \xrightarrow{\alpha} z$$

in the tree $\overline{Q}$. In particular, we assume that $d(x, z) = d(x, y) + 1$. First let us stress a general fact which does not depend on the orientation of α .

(0) *Any vertex in $\mathcal{B}_z$ is connected to precisely one vertex in $\mathcal{B}_y$.*

Proof. This is an immediate consequence of the way the radiation quiver is constructed. Namely, any vertex b of $R(M, x)$ occurs at a certain step as an origin. In the next step, this vertex b is connected by an arrow to the new origin, say b' . In later steps, no further arrows are attached which involve b . Thus, let $d(x, z) = i$ and let D be the maximal distance of vertices to x . We start with step 0, namely with the disjoint union of radiation quivers consisting of a single vertex, one for each vertex with distance D to x . The basis $\mathcal{B}_z$ is constructed in step $D - i$, all the elements of $\mathcal{B}_z$ are origins of radiation quivers. In the step $D - i + 1$, the basis $\mathcal{B}_y$ is constructed: here, any vertex $b \in \mathcal{B}_z$ is connected by an edge to precisely one vertex $b' \in \mathcal{B}_y$.

Corollary. *The number of edges labeled α is equal to $\dim M_z$.*

Proof: By (0), the number of these edges is equal to the cardinality of $\mathcal{B}_z$, but this is just $\dim M_z$.

Now we take into account the orientation of α . Actually, we distinguish four cases: the arrow α may point into two different directions and the corresponding map may be injective or surjective.

(1) *If $\alpha: y \rightarrow z$ is injective, then any vertex in $\mathcal{B}_y$ is connected by an edge to at least one vertex in $\mathcal{B}_z$.*

Proof: If $b \in \mathcal{B}_y$ is not connected by an edge to a vertex in $\mathcal{B}_x$, then b belongs to the kernel of M_α .

(2) If $\alpha: y \rightarrow z$ is surjective, then any vertex in $\mathcal{B}_y$ is connected by an edge to at most one vertex in $\mathcal{B}_z$.

Proof: If there are vertices $b' \neq b''$ in $\mathcal{B}_z$ which are connected by edges to $b \in \mathcal{B}_y$, then neither b' nor b'' belong to the image of M_α .


(3) If $\alpha: z \rightarrow y$ is injective, then any vertex in $\mathcal{B}_y$ is connected by an edge to at most one vertex in $\mathcal{B}_z$.

Proof: If there are vertices $b' \neq b''$ in $\mathcal{B}_z$ which are connected by edges to $b \in \mathcal{B}_y$, then $b' - b''$ belongs to the kernel of M_α .

(4) If $\alpha: z \rightarrow y$ is surjective, then any vertex in $\mathcal{B}_y$ is connected by an edge to at least one vertex in $\mathcal{B}_z$.

Proof: If $b \in \mathcal{B}_y$ is not connected by an edge to a vertex in $\mathcal{B}_z$, then b is not in the image of M_α .

It follows that the edges between $\mathcal{B}_y$ and $\mathcal{B}_z$ are of one of the following form:



3. Dynkin quivers.

Proposition 3. Let Q be a Dynkin quiver and M an indecomposable thin vertex x . Then (M, x) is a radiation module.

This is an immediate consequence of the following lemma.

Lemma 3. Let Q be a Dynkin quiver and M an indecomposable with thin vertex x . Let M' be the restriction of M to the quiver obtained by deleting the vertex x . Then M' decomposes as the direct sum of pairwise non-isomorphic indecomposable modules $N(i)$. For every index i , there is a neighbor $y(i)$ of x with $\dim N(i)_{y(i)} = 1$ and this is the unique neighbor y of x such that $\dim N(i)_y \neq 0$.

Proof of lemma 3. For a Dynkin quiver, all indecomposable representations are exceptional, thus we can apply proposition 1 and decompose $M' = \bigoplus N(i)^{n(i)}$ where

$N(1), \dots, N(t)$ is an orthogonal exceptional family. If $n(i) \geq 2$ for some i , then the process of simplification shows that the full subcategory of all modules with a filtration with factors $S(x)$ and $N(i)$ is representation infinite, impossible. Thus $M' = \bigoplus N(i)$ with pairwise non-isomorphic indecomposable modules $N(i)$. As proposition 1 asserts, for every i , there is a unique vertex y with $\dim N(i)_y \neq 0$ and $\dim N(i)_{y(i)} = n(i) = 1$.

Note that for Δ a Dynkin quiver, the only exceptional modules without a thin vertex are the maximal indecomposable modules for the quivers of type $\mathbb{E}_8$.

4. Bipartite trees without leaves

Let us now draw the attention to a class of infinite quivers, but we will assume that the quivers are locally finite. For a general discussion of the Auslander-Reiten components of infinite quiver we may refer to [BLP].

The quiver Q is said to be *bipartite* provided that every vertex is a sink or a source. A bipartite tree quiver will just be called a *bipartite tree*. Note that any graph which is a tree can be endowed with precisely two orientations so that we obtain a bipartite tree. A vertex x of a tree quiver is called a *leaf* provided there is at most one arrow α which starts or ends in x . Note that a tree quiver without leafs has to be infinite. As typical examples one should take an n -regular tree with bipartite orientation, where $n \geq 2$. These are the quivers which we later will use when dealing with the generalized Kronecker quivers. We should stress that the case $n = 3$ was already discussed in [FR].

If Q is a quiver, let Q^* be the opposite quiver: it has the same vertices, but every arrow $\alpha: x \rightarrow y$ is replaced by an arrow $\alpha^*: y \rightarrow x$.

Let Q be a bipartite tree. Given two vertices x, y , let $d(x, y)$ be the distance between x and y . We denote by $N(x)$ the set of neighbors of x (thus the set of vertices y with $d(x, y) = 1$).

If $x \in Q_0, t \in \mathbb{N}_0$, let $B(x, t)$ be the set of vertices y with $d(x, y) \leq t$, we may call it the *ball* with center x and radius t . If necessary, we will consider $B(x, t)$ as a full subquiver of Q . We will be interested in the balls $B(x, t)$ where x is a sink and t is even, as well in those where x is a source and t is odd. Note that for these pairs x, t , the boundary $\{y \mid d(x, y) = t\}$ of $B(x, y)$ consists of sinks (if x is a sink, then y is also a sink iff $d(x, y)$ is even. If x is a source, then y is a sink iff $d(x, y)$ is odd).

We denote by ρ^- the composition of all BGP-reflection functors at all the sources of Q , this is a functor $\rho^-: \text{mod } kQ \rightarrow \text{mod } kQ^*$. Note that $\rho^- \rho^-$ is just the Auslander-Reiten translation τ^- , see Gabriel [G].

Given a bipartite quiver Q without leaves, let us define indecomposable representations $P(x, t) = P_Q(x, t)$ for all $x \in Q_0, t \in \mathbb{N}_0$ as follows: If x is a sink, let $P(x, 0) = S(x) = P(x)$, the one-dimensional representation with support $\{x\}$. If x is a source, let $P(x, 1) = P(x)$, the indecomposable projective module with top $S(x)$; it is the unique thin module with support $B(x, 1)$. For $t \geq 2$, define $P(x, t) = \tau^- P(x, t-2)$. Thus, by definition

$$\begin{aligned} P(x, 2t) &= \tau^{-t} P(x, 0), \\ P(x, 2t+1) &= \tau^{-t} P(x, 1). \end{aligned}$$

Looking at the same time at the quivers Q and Q^* , we see that for all $t \geq 1$

$$P_Q(x, t) = \rho^- P_{Q^*}(x, t - 1).$$

We call the modules of the form $P(x, t)$ the preprojective modules.

Lemma 4.

- (a) *The support of $P(x, t)$ is $B(x, t)$.*
- (b) *If $t \geq 1$ and $d(x, y) \in \{t - 1, t\}$, then $\dim P(x, t)_y = 1$.*
- (c) *Let X be an indecomposable representation of Q . Then $\text{Hom}(X, P(x, t)) \neq 0$ if and only if X is preprojective and the support of X is contained in the support of $P(x, t)$.*
- (d) *Let $P = P(x, t)$ and z a vertex with $P_z = 0$. Then $d(x, z) = t + 1$ if and only if $\text{Ext}^1(S(z), P) \neq 0$ and then $\dim \text{Ext}^1(S(z), P) = \dim P_y = 1$.*

Proof: (b) We use induction, the case $t = 0$ being trivial. Let $t \geq 1$. We know that $P_Q(x, t) = \rho^- P_{Q^*}(x, t - 1)$. Let y be a vertex of Q with $d(x, y) = t$. There is a unique neighbor y' of y with $d(x, y') = t - 1$ and $\dim P_{Q^*}(x, t - 1)_{y'} = 1$ (this is clear for $t = 1$, whereas for $t > 1$ this is the induction hypothesis). The definition of ρ^- shows that $P(x, t)_y = P_{Q^*}(x, t - 1)_{y'}$, thus $\dim P(x, t)_y = 1$. If $t \geq 1$ and y' is any vertex of Q with $d(x, y') = t - 1$, then ρ^- does not change the vector space at the position y' , thus $\dim P(x, t)_{y'} = \dim P_{Q^*}(x, t - 1)_{y'} = 1$.

(a) The induction procedure mentioned in (a) shows that the support of $P(x, t)$ is contained in $B(x, t)$, and as we have seen, it contains all the vertices y with $d(x, y) = 1$. But the support of an indecomposable module has to be connected, thus it has to be all of $B(x, t)$.

(c) Let X be an indecomposable representation of Q with $\text{Hom}(X, P(x, t)) \neq 0$. We write $P(x, t) = \tau^{-s} P$ for some indecomposable projective module (namely $P = \tau^s P(x, t)$ and $s = \lfloor \frac{t}{2} \rfloor$). If $\tau^s X \neq 0$, then $\text{Hom}(\tau^s X, P) \neq 0$ and therefore $\tau^s X$ is projective. Altogether, we see that $X = \tau^{-i} P'$ for some indecomposable projective module $P' = P(y)$ and $0 \leq i \leq s$. Since $\text{Hom}(P', \tau^{-s+i} P) = \text{Hom}(P', \tau^i P(x, t)) \neq 0$, it follows that the y belongs to the support of $\tau^i P(x, t)$, thus to the support $B(x, t)$ of $P(x, t)$.

Conversely, consider the quiver $B(x, t)$. Since this is a tree quiver, the category of representations of $B(x, t)$ has a preprojective component. Let $\mathcal{C}(x, t)$ be the full subcategory of the indecomposable representations of $B(x, t)$ which are predecessors of $P(x, t)$, we may consider this as a subcategory of the category of representations of Q . As we have seen, any predecessor X of $P(x, t)$ in the category $\text{mod } kQ$ belongs to $\mathcal{C}(x, t)$.

(d) Let $P = P(x, t)$ and z a vertex with $P_z = 0$ and $\text{Ext}^1(S(z), P) \neq 0$. Since $P_z = 0$, we must have $d(x, z) > t$. Since $\text{Ext}^1(S(z), P) \neq 0$, we see that $d(x, z) = t + 1$. Then $\dim \text{Ext}^1(S(z), P) = \dim P_y = 1$, according to (b).

Proposition 4. *The preprojective modules are reachable and they form a component of the Auslander-Reiten quiver.*

Here is a description of the Auslander-Reiten sequences starting in $P(x, t)$


$$0 \rightarrow P(x, t) \rightarrow \bigoplus_{y \in N(x)} P(y, t + 1) \rightarrow P(x, t + 2) \rightarrow 0.$$

It is of interest to stress the following property of this preprojective component $\mathcal{P}$: for any indecomposable object X in $\mathcal{P}$, there are sectional paths starting in a simple projective module S and ending in X , namely, for $X = P(x, t)$ there are sectional paths from $S = P(y, 0)$, where $d(x, y) = t$.

Proposition 5. *For $t \geq 1$, the pairs $(P(x, t), y)$ with $d(x, y) \in \{t, t-1\}$ are radiation modules.*

Proof. It is sufficient to consider a vertex y with $d(x, y) = t-1$. We know that $\dim P(x, t)_y = 1$. Let U be the submodule of $P(x, t)$ with $P(x, t)/U = S(y)$. Since U is a submodule of $P(x, t)$, we know that U is a direct sum of preprojective modules P , and any such P satisfies $P_y = 0$ and $\text{Ext}^1(S(y), P) \neq 0$. According to (d), we know that $\dim \text{Ext}^1(S(y), P) = 1$. Now use proposition 1.

Centipedes. Given a pair x, y of vertices of Q , we denote by $[x, y]$ the set of vertices lying on the path between x and y . If S is a set of vertices of Q , we denote by $N(S)$ the set of neighbors of S : these are the vertices z which do not belong to S but such that there is a vertex $z' \in S$ with $d(z, z') = 1$. We will be interested in the sets $N([x, y])$, where x, y are vertices with y a source:



Lemma 5. *Let y be a source and $z \neq y$ some other vertex. Then $S(y), P(z, d(y, z)-1)$ is an orthogonal exceptional pair with $\dim \text{Ext}^1(S(y), P(z, d(y, z)-1)) = 1$.*

Proof. According to (a), we have $P(z, d(y, z)-1)_y = 0$, thus $S(y), P(z, d(y, z)-1)$ is an orthogonal pair. And this pair is exceptional, since $S(y)$ is injective. According to (d), we have $\dim \text{Ext}^1(S(y), P(z, d(y, z)-1)) = 1$.

Proposition 6. *Let y be a source and x any vertex. Let $d(x, y) = t-1$ for some $t \geq 1$. Then there is the following exact sequence:*

$$0 \rightarrow \bigoplus_{z \in N([x, y])} P(z, t-d(x, z)) \rightarrow P(x, t) \rightarrow S(y) \rightarrow 0$$

The modules $S(y)$ and $P(z, t-d(x, z))$ with $z \in N([x, y])$ are pairwise orthogonal bricks and the map $P(x, t) \rightarrow S(y)$ is the projective cover of $S(y)$ in the full subcategory

$$\mathcal{F}\left(S(y); P(z, t-d(x, z)), z \in N([x, y])\right).$$

Proof. We use induction on t . In the case $t = 1$, we have $x = y$ and there is the exact sequence

$$0 \rightarrow \bigoplus_{z \in N(x)} P(z, 0) \rightarrow P(x, 1) \rightarrow S(x) \rightarrow 0.$$

Now assume the assertion is true for some $t \geq 1$. Consider a pair of vertices x, z_0 with $d(x, z_0) = t$. There is a unique vertex y with $d(x, y) = t - 1$ and $d(y, z_0) = 1$. By induction, there is the exact sequence

$$(*) \quad 0 \rightarrow \bigoplus_{z \in N([x, y])} P(z, t - d(x, z)) \rightarrow P(x, t) \rightarrow S(y) \rightarrow 0.$$

Choose some neighbor z_0 of y with $d(x, z_0) = t$. There is (up to isomorphism) a unique module M with top $S(y)$ and socle $S(z_0)$ and we may rearrange the factors in $(*)$ in order to obtain an exact sequence of the form

$$0 \rightarrow \bigoplus_{z \in N([x, y]), z \neq z_0} P(z, t - d(x, z)) \rightarrow P(x, t) \rightarrow M \rightarrow 0.$$

Now apply the functor ρ^- . We obtain the sequence

$$(**) \quad 0 \rightarrow \bigoplus_{z \in N([x, y]), z \neq z_0} P(z, t + 1 - d(x, z)) \rightarrow P(x, t + 1) \rightarrow \rho^- M \rightarrow 0.$$

The module $\rho^- M$ has top $S(z_0)$ and its socle is the direct sum of the modules $S(z)$ where $z \neq y$ is a neighbor of z_0 . Of course, for these vertices z , we have $S(z) = P(z, 0)$ and $d(x, z) = t + 1$. Since these modules $P(z, 0)$ are projective, we obtain from $(**)$ an exact sequence of the form


$$0 \rightarrow U \rightarrow P(x, t + 1) \rightarrow S(z_0) \rightarrow 0$$

where U is the direct sum of the modules $P(z, t + 1 - d(x, z))$ with $z \in N([x, y]), z \neq z_0$ as well as those of the form $P(z, 0) = P(z, t + 1 - d(x, z))$ with $z \neq y$ a neighbor of z_0 . It remains to observe that the elements in $N([x, z_0])$ are precisely the vertices $z \in N([x, y]), z \neq z_0$ and the neighbors $z \neq y$ of z_0 . Thus we see that we obtain for the pair x, z_0 the required sequence.

Now let us show that the modules $S(y)$ and $P(z, t - d(x, z))$ and $z \in N([x, y])$ are orthogonal and that the only non-trivial extension groups between these modules are the groups $\text{Ext}^1(S(y), P(z, t - d(x, z)))$ and these are one-dimensional. We can assume that $t \geq 2$ and we denote by y' the unique vertex with $d(y, y') = 1$ and $d(y', x) = t - 2$. If $d(x, z) = t$, then $P(z, t - d(x, z)) = P(z, 0) = S(z)$ is simple projective, not isomorphic to $S(y)$ and $\dim \text{Ext}^1(S(y), S(z)) = 1$. Also, for $z \neq z'$, the modules $S(z), S(z')$ are orthogonal. Thus, assume that $d(x, z) < t$. Let $z' \in [x, y]$ with $d(z, z') = 1$. Write $a = d(y, z')$ and $b = d(z', x)$, thus $a + b = t - 1$ and $d(z, x) = b + 1$, thus $t - d(x, z) = a$. Since $d(x, z) < t$, we see that $z' \neq y$, thus $d(z, y') = a$ and therefore $P(z, a)_y = 0$ and $\dim P(z, a)_a = 1$. According to (d), we have $\dim \text{Ext}^1(S(y), P(z, a)) = 1$. It follows from $P(z, a)_y = 0$ that $P(z, a)_{z'} = 0$ for $d(x, z') = t$, thus $P(z, a)$ and $P(z', 0)$ are orthogonal, and also that $\text{Ext}^1(P(z, a), P(z', 0)) = 0$. Assume now that we deal with two vertices $z_1 \neq z_2$ in $N([x, y])$ with $d(x, z_2) \leq d(x, z_1) < t$, let $a_1 = t - d(x, z_1)$, and $a_2 = t - d(x, z_2)$, thus $a_1 \leq a_2$. In order to see that there are no homomorphisms or extensions between $P(z_1, a_1)$ and $P(z_2, a_2)$, we can apply ρ^{a_1} , thus we have to consider instead $P(z_1, 0)$ and $P(z_2, a_2 - a_1)$. Note that $d(z_1, z_2) = a_2 - a_1 + 2$, thus z_1 is not in the support of $P(z_2, a_2 - a_1)$ and not even a neighbor of this support. This completes the proof.


Corollary. *Let y be a source and x any vertex. The family of modules $P(z, d(y, z) - 1)$ with $z \in N([x, y])$ is an orthogonal family of bricks without mutual extensions.*

As an example, let us consider the 3-regular tree with bipartite orientation as studied already in [FR]. Let us display the dimension vector $\mathbf{dim} P(x, 3)$ for a source x . As vertex y we take a source with $P(x, 3)_y$ being one-dimensional, so that $(P(x, 3), y)$ is a radiation module.




Here, x is the vertex at the center. The vertex y is encircled, using a solid circle, the vertices $z \in N([x, y])$ are encircled using dashed circles.

Here is the radiation quiver $R(P(x, 3), y)$.



Finally, let us show the dimension vector $\mathbf{dim} P(x, 4)$ with x a sink. Again, we choose a source y such that $\dim P(x, 4)_y = 1$.



Again, x is the vertex at the center. The vertex y is encircled, using a solid circle, whereas the dashed circles mark the vertices $z \in N([x, y])$. Altogether, we see that there is an exact sequence of the form

$$0 \rightarrow U \rightarrow P(x, 4) \rightarrow S(y) \rightarrow 0$$

with

$$U = P(z_1, 0) \oplus P(z_2, 0) \oplus P(z_3, 1) \oplus P(z_4, 2) \oplus P(z_5, 3) \oplus P(z_6, 3)$$

6. The generalized Kronecker quivers and Schofield induction.

The generalized Kronecker quivers are the quivers $K(n)$ with 2 vertices, a sink and a source, and n arrows (going from the source to the sink); the case $n = 2$ is the ordinary Kronecker quiver, its representations have been studied by Weierstraß and Kronecker. The universal covering $Q(n)$ of the quiver $K(n)$ is the n -regular tree with bipartite orientation. Using the push-down functor, any indecomposable representation of the quiver $Q(n)$ yields an indecomposable representation of $K(n)$. In particular, the representation $P_{Q(n)}(x, t)$ of $Q(n)$ defined in the last section provides an indecomposable representation $P_{K(n)}(t)$ and one obtains in this way just the preprojective $K(n)$ -modules (the special case $n = 3$ has been discussed in detail in [FR], all the considerations presented there can easily be adapted to the general case). Obviously, under the push-down functor a tree basis is sent to a tree basis. Since all the modules $P(x, t)$ are radiation modules we obtain in this way

distinguished tree bases for the preprojective $K(n)$ -modules. The dual considerations yield distinguished tree bases for the preinjective $K(n)$ -modules.

Let us consider now an arbitrary quiver Q and M an exceptional representations of Q . It has been shown in [R2] (see also [R3]) that M is a tree module. Here we want to outline in which way we can use the previous assertions in order to exhibit a nice tree basis of M . In order to exhibit a tree basis for an exceptional module M , one uses Schofield induction, thus one considers exact sequences of the form

$$0 \rightarrow Y^y \rightarrow M \rightarrow X^x \rightarrow 0$$

with indecomposable middle term, where (X, Y) is an orthogonal exceptional pair with $\dim \text{Ext}^1(X, Y) = e > 0$ and (x, y) is the dimension vector of a sincere preprojective or preinjective representation E of the e -Kronecker quiver $K(e)$. Note that the triple $(X, Y; E)$ uniquely determines M and one obtains in this way inductively all the exceptional representations of Q . In order to construct a tree basis of M we need to know tree bases of X, Y and E . For the procedure to obtain a tree basis of M from the tree bases of X, Y, E we refer to [R2] (see also [R3]).

A required tree basis for E has been exhibited already in [R2], but the present note provides an intrinsic way for obtaining such a tree basis. Let us repeat: Instead of working with the e -Kronecker quiver itself, we consider its universal cover, the e -regular tree $Q(e)$ with bipartite orientation. Let E be obtained from the representation $\tilde{E}$ of $Q(e)$ by the push-down functor. Now $Q(e)$ is a bipartite tree without leaves and $\tilde{E}$ is a preprojective or preinjective representation of Q . Thus, there is a vertex x such that the pair $(\tilde{E}, x)$ is a radiation quiver. But this means that $\tilde{E}$ has a distinguished tree basis. Under the push-down functor, we obtain a distinguished tree basis for E .

References

- [BLP] R. Bautista, S. Liu, and C. Paquette, Representation theory of strongly locally finite quivers, Proc. London Math. Soc. (to appear).
- [FR] Ph. Fahr, C. M. Ringel: Categorification of the Fibonacci numbers using representations of quivers. Journal of Integer Sequences. Vol. 15 (2012), Article 12.2.1
- [G] P. Gabriel: Auslander-Reiten sequences and representation-finite algebras. In Representation theory I. Springer LNM 831, Berlin (1980) 171.
- [R1] C. M. Ringel: Tame algebras and integral quadratic forms. Springer LNM 1099 (1984).
- [R2] C. M. Ringel: Exceptional modules are tree modules. Lin. Alg. Appl. 275-276 (1998) 471-493.
- [R3] C. M. Ringel: Indecomposable representations of the Kronecker quivers. Proceedings AMS (to appear).

C. M. Ringel
 Fakultät für Mathematik,
 Universität Bielefeld, PO Box 100 131
 D-33501 Bielefeld, Germany
 e-mail: ringel@math.uni-bielefeld.de